

STIC-ILL

462081

ND

From: Canella, Karen  
Sent: Tuesday, September 02, 2003 2:46 PM  
To: STIC-ILL  
Subject: ill order 09/541,345

Art Unit 1642 Location 8E12(mail)

Telephone Number 308-8362

Application Number 09/541,345

1. Semin Cancer Biol, 1999 Apr, 9(2):67-81
2. Int J Cancer, 1998 Mar 30, 76(1):148-153
3. Int J Biol Markers:  
1992 Jul-Sep, 7(3):137-142  
1992 Jul-Sep, 7(3):132-136
4. Gene, 1993, 128(1):29-36

11551692

# The CEA family: *a system in transitional evolution?*

C. P. STANNERS, M. ROJAS, H. ZHOU, A. FUKS, N. BEAUCHEMIN

McGill Cancer Centre and Departments of Biochemistry and Oncology, McGill University, Montreal, Quebec-Canada

**ABSTRACT:** The CEA family consists of two structurally and functionally distinct sub-groups; the group including CEA, NCA and CGM-6 which are cell surface-bound by phosphatidyl-inositol (PI) linkages, and the group of BGP splice variants which have trans-membrane and cytoplasmic domains. Although all CEA family members mediate intercellular adhesion *in vitro*, the PI-linked group show  $Ca^{++}$  and temperature independent adhesion whereas the BGP group show rapidly reversible  $Ca^{++}$  and temperature dependent adhesion. From the close alignment in cDNA nucleotide sequences between family members and between repeated domains in one family member, it is apparent that the CEA family is now rapidly evolving; in fact, analogs of only the trans-membrane BGP group have been found so far in the mouse. The addition of a new group of potent adhesion molecules to complex species at some time after the rodent radiation has strong evolutionary implications, which are discussed. (*Int J Biol Markers*, 1992; 7: 137-42)

**KEY WORDS:** CEA family, Intercellular adhesion, CEA function

## INTRODUCTION

Human carcinoembryonic antigen (CEA) is a member of a family of cell surface glycoproteins which appear in the blood in association with a wide variety of cancers, including colon, breast and lung (1). The cloning of genes and cDNA's for CEA in 1987 (2-8) allowed the delineation of the complete structure of CEA itself and a means for clarification of the extent of the family and the precise structures of all of its members. Considering only the CEA branch of the CEA family (the other branch being the pregnancy-specific glycoproteins), 9 genes have been identified which are closely clustered in two groups on human chromosome 19 (9, 10, 11). Figure 1 depicts the basic structures of the CEA family members for which protein products have been reasonably well characterized. As a subset of the immunoglobulin supergene family, all members consist of a processed leader sequence, followed by a V-like N-terminal domain, a variable number (from 0 to 6) of C2-set domains each with a disulphide bridge, and either a hydrophobic C-terminal domain, which is processed to give a phosphatidyl-inositol (PI) type of linkage to the external cell membrane (CEA, NCA and HCGM-6) (14-17), or a transmembrane domain with a short or a long cytoplasmic tail (splice variants of BGP) (18, 19).

The function(s) of the CEA family was completely unknown prior to the availability of full-length cDNA

clones which allowed functional studies in transfectants. CEA (20, 21), NCA (21, 22, 24) and BGP (23) have all been shown to function *in vitro* as homotypic intercellular adhesion molecules using stable transfectant clones of previously non-adhesive rodent cell lines. HCGM-6 transfectants, on the other hand, appear to interact only heterotypically with NCA transfectants (22, Ilyantis and Stanners, unpublished results). These data together with results showing a lack of heterotypic interactions with transfectants of other adhesion molecules of the immunoglobulin supergene family such as N-CAM and L-MAG (24), indicate that the interactions are specific.

The characteristics of the adhesive process for the phosphatidyl-inositol linked family members (CEA, NCA and HCGM-6) differ from those of the transmembrane linked family members (BGP) in that the BGP members require calcium and physiological temperatures for adhesion while the CEA group show adhesive function in the absence of calcium and at 4° C (20-23).

In addition to this intercellular adhesive function for CEA family members, the sugars of CEA, especially the mannose-rich sugar structures on one of the glycoforms of NCA, have been shown to bind gram negative bacteria (25). The question of further functions for CEA remains open, especially since other receptors which bind CEA have been demonstrated (26).

## CEA FAMILY




Fig. 1 - Structure of several members of the CEA family. The V-like Ig N domains are shown as linear termini, while the C2-set domains are shown with loops linked by disulphide bridges. PI-linkage to the external cell membrane is depicted with an arrowhead; transmembrane and cytoplasmic domains are shown for the BGP members, although the configuration with respect to the cell membrane for BGP<sub>g</sub> and BGP<sub>h</sub> is inferred from the nucleotide-amino acid sequence.

Notwithstanding this recent progress in delineating *in vitro* functions for CEA, the demonstration that these function(s) apply *in vivo* has yet to be shown. This, of course, represents a very difficult task. If the intercellular adhesion function also applies *in vivo*, one might expect a role in embryonic development and/or in the maintenance of tissue architecture. It is the purpose of this article to use comparative data from both human and animal systems in order to examine the question of function from an evolutionary perspective.

## RESULTS

**Further delineation of BGP function(s):** the calcium and temperature dependence observed for intercellular adhesion mediated by BGP<sub>a</sub> and BGP<sub>b</sub> (23) represent important attributes of adhesion which are also characteristic of the cadherin family of adhesion molecules (27), although, recently, calcium-dependent adhesion has been observed for another member of the immunoglobulin supergene family, PECAM-1 (28). This does not imply that the BGP members are necessarily involved in the same types of functions as the cadherins, such as the initiation and maintenance of functional intercellular junctions. It does, however, raise the possibility that the BGP-mediated adhesion

could be modulated rapidly by changes in extracellular conditions such as divalent cation concentrations, perhaps by direct "ecto" effects involving the extracellular domains of the molecules. These extracellular events could include direct effects of divalent cations on molecular conformation, ecto-ATPase activity (29), and ecto-phosphorylation of BGP molecules, as suggested by Obrink for the rat analog of human BGP, cell-CAM 105 (30). It is of interest that a cDNA cloned using amino-acid sequence information derived from purified rat ecto-ATPase turned out to be identical to this relatively well characterized adhesion molecule (31). We have looked for ecto-ATPase activity in purified preparations of human BGP<sub>a</sub> using several assays, without convincingly positive results (Sadoul-Schellenberger and Stanners, unpublished results); it is possible, however, that such activity could depend on integration of BGP molecules in the cell membrane or on associated molecules. We have shown, nevertheless, that BGP adhesion depends on Mg<sup>++</sup> as well as Ca<sup>++</sup> (data not shown). This makes it more likely that the effect of these cations is direct, presumably on molecular conformation, i.e., "ecto", rather than indirect through effects on the physiological state of the cell. Work is in progress to study the effect of divalent cations on the molecular conformation of BGP directly.

With regard to reversibility of adhesion, the aggregation of BGP transfectants has been studied as a function of changing extracellular conditions. For these studies, clones of the CHO derived cell line, LR-73 (32), stably transfected with functional cDNA of a splice variant of BGP known as BGP<sub>h</sub>, were used. This variant contains the least of the possible coding sequences found in BGP splice variants, consisting of the leader, the first 68 amino acids of the N domain coupled directly to the transmembrane domain and the short cytoplasmic domain of 9 amino acids (T. Barnett, personal communication). Adhesion was studied, in comparison with stable LR-73 transfectants of CEA and E-cadherin, as a function of temperature (Fig. 2) and extracellular ATP (Fig. 3). In the first place it is evident that this severely truncated BGP variant is capable of mediating adhesion, unlike CEA whose complete N domain alone is incapable of adhesion (Zhou and Stanners, in preparation). Secondly, the cellular aggregates formed through the action of this molecule rapidly dissociate in response to a temperature shift from 37°C to 4°C, and rapidly reform when the temperature is shifted back to 37°C (Fig. 2). These changes were not observed for transfectants of CEA (Fig. 2) or for NCA transfectants (data not shown). In




Fig. 2 - Aggregation of BGP<sub>b</sub> and CEA transfectants as a function of rapid shifts in temperature. Clones of LR-73 cells stably transfected with functional CEA or BGP<sub>b</sub> cDNA were produced and grown in monolayer culture as previously described (20). After rendering cultures single cell suspensions, the percentage of single cells during the formation and dissociation of aggregates was measured with time as previously described (20). Rapid changes in temperature were effected at the indicated times.

addition, aggregates of BGP transfectant cells dissociated immediately after the addition of ATP, whereas CEA and E-cadherin transfectants were unaffected (Fig. 3). This effect was not seen with ADP (data not shown), and could be due to ATP forming complexes and removing  $\text{Ca}^{++}$  from the medium or could be a direct effect through an ecto-ATPase or ecto-phosphorylation function. Similar rapid reversibility of aggregation was obtained with adjustment of the  $\text{Ca}^{++}$  concentration in the cellular suspension medium (data not shown).

**Extent of CEA family in mice:** a restriction fragment of human CEA cDNA from the translational start codon to about the middle of the second C2-set repeated domain was used as a probe to isolate cDNA clones encoding mouse analogs of the human CEA family. A series of clones representing splice variants of what is probably a single gene have been characterized (33, 34). These all have transmembrane domains and




Fig. 3 - Aggregation of BGP<sub>b</sub>, CEA, and E-cadherin transfectants before and after the addition of ATP. The aggregation and dissociation with time of suspended cells from cultures of stable BGP<sub>b</sub>, CEA and E-cadherin cDNA transfectant clones in the absence and presence of 1mM ATP were measured as described in the legend of Figure 1.

either a short or a long cytoplasmic domain of 10 or 73 amino acids, with variable N and C2-set domains. Transfectants of mouse cells with two of these murine splice variants have been shown to demonstrate homotypic intercellular adhesion (33,34) and one has been shown to be temperature and  $\text{Ca}^{++}$ -dependent (33). The other effected  $\text{Ca}^{++}$ -independent adhesion, but this could have been due to a very high level of production observed for this particular transfectant clone, an effect also observed for high producing transfectants of human BGP (Rojas and Stanners, in preparation). In addition, the tissue-specific pattern of expression of the murine analogs resembles that of human BGP (30, 33). It should also be pointed out that the amino-acid sequence alignment of the murine family members is closer with human BGP than with human CEA or NCA (33, 34). Thus, all of the murine CEA-like cDNA clones isolated to date resemble human BGP both in structure, function and expression patterns. No transcripts corresponding to PI-linked murine family mem-

bers analogous to CEA, NCA and HCGM-6 have been discovered to date either in cDNA libraries of murine colon, on Northern blots of RNA from many murine tissues, or by PCR amplification of mouse colon and liver RNA (McCuaig et al, in preparation).

## DISCUSSION

The above results add to our previous data showing that the BGP members of the human CEA family differ markedly in the characteristics of their intercellular adhesion function from those of the PI-linked members. These differences include  $\text{Ca}^{++}$  dependence, temperature dependence and immediate reversibility of adhesion by extracellular conditions; the latter imply "ecto" control of adhesion through rapid changes in molecular conformation. Where tested, these functional properties are shared by the CEA-like molecules which we have detected in the mouse. This leads to the suggestion that the PI-linked members of the human CEA family represent an evolution of a new branch of the family with an evolved adhesive function.

Although it may be presumptuous to suggest that the PI-linked CEA family members do not in fact exist in the mouse, all our evidence to date points in that direction; definitive evidence will require detailed genomic studies and/or careful PCR analysis of mRNA from many tissues of the mouse. If so, we are faced with the prospect of a group of fairly potent intercellular adhesion molecules which have been introduced into the human lineage at some point after the radiation of murine species. This suggestion of a recent evolution of the CEA family is supported by the very close nucleotide sequence alignment (> 80%) between the repeated double C2-set domains in CEA and between CEA, NCA and HCGM-6. Of further interest is the fact that, although there are regions where very little change has occurred, the amino-acid sequence alignment in these comparisons is always less than the nucleotide sequence alignment (4). This is, in fact, exactly as predicted for random changes in the nucleotide sequence without preservation of the actual amino-acid sequence (4). What has been conserved is, as mentioned above, certain limited regions of both the nucleotide and the amino-acid sequence, and the open reading frames; what has evolved are changes in the carboxy terminal sequence to effect PI-linkage. It is possibly the latter linkage that has allowed the evolution of intercellular adhesion molecules with different adhesive characteristics.

The PI-linked CEA family molecules must present

some advantage to the organism as they have evolved as such and their open reading frames have been conserved. If they do, in fact, function as intercellular adhesion molecules during embryonic development and in adult tissues, how can mice with their very similar developmental and adult tissue architecture do without them (assuming that the failure to detect them in mice is confirmed)? The answer is, perhaps, a testament to the plasticity of complex biological systems. Since many genes can be "knocked out" in such organisms without apparent phenotypic effect, it is clear that conserved genes with advantageous functions must exist and persist by conferring an inapparent selective advantage. Consistent with this view, the addition of functional CEA nucleotide sequences driven by the SV40 transcriptional promoter to the genome of mice by transgenesis had no apparent effect on their phenotype, even though expression was observed in many tissues (35).

These considerations may provide an explanation for the paradox of the existence of a group of genes in humans but not in mice with a potential function which should affect development. They do not, of course, ratify an intercellular adhesion function for these genes *in vivo*. A highly speculative suggestion is that the PI-linked members of the CEA family represent part of a gene family in "transitional evolution". A simplistic view of molecular evolution would envision the replication of existing genes or domains/exons followed by drift in the amino-acid sequence until a new or modified function capable of conferring selective advantage emerges. All that is required in the transitional period are preservation of the open reading frame and a function which has small advantage and which is at least not deleterious. The CEA family in man could presently be in such a state as viewed through the present narrow window in evolutionary time. The significance of overproduction of the PI-linked family members in tumours is, of course, a completely separate issue.

## ACKNOWLEDGEMENTS

We thank Dr. T. Barnett for providing a cDNA clone corresponding to BGP<sub>h</sub>.

This work was supported by a program grant from the Medical Research Council of Canada to C.P.S., A.F. and N.B. and by grants from the N.C.I.(C) to C.P.S. and A.F.

Reprint requests to:  
C.P. Stanners, MD  
McGill Cancer Centre and  
Departments of Biochemistry and Oncology,  
McGill University,  
3655 Drummond St.  
Montreal, Quebec  
Canada H3G1Y6

## REFERENCES

- Shuster JD, Thompson MP, Fuks A, Gold P. Immunologic approaches to diagnostic of malignancies. *Prog Exp Tumor Res* 1980; 25: 89-139.
- Schrewe H, Thompson J, Bona M, Hefta LJF, Maruya A, Hassauer M, Shively JE, von Kleist S, Zimmermann W. Cloning of the complete gene for carcinoembryonic antigen: analysis of its promoter indicates a region conveying cell type - specific expression. *Mol Cell Biol* 1990; 10: 2748.
- Willcocks TC, Craig IW. Characterization of genomic organization of human carcinoembryonic antigen (CEA): comparison with other family members and sequence analysis of 5' controlling region. *Genomics* 1990; 8: 492-500.
- Beauchemin N, Benchimol S, Cournoyer D, Fuks A, Stanners CP. Isolation and characterization of full-length functional cDNA clones for human carcinoembryonic antigen. *Mol Cell Biol* 1987; 7: 3221-30.
- Oikawa S, Nakazato H, Kosaki G. Primary structure of human carcinoembryonic antigen (CEA) deduced from cDNA sequence. *Biochem Biophys Res Comm* 1987; 14: 511-8.
- Zimmermann W, Ortlieb B, Friedrich R, von Kleist S. Isolation and characterization of cDNA clones encoding the human carcinoembryonic antigen reveal a highly conserved repeating structure. *Proc Natl Acad Sci USA* 1987; 84: 2960-4.
- Kamarck ME, Elting JJ, Hart JT, Goebnel SJ, Rac PMM, Nothdurft MA, Nedwin JJ, Barnett TR. Carcinoembryonic antigen family: expression in a mouse L-cell transfectant and characterization of a partial cDNA in bacteriophage lambda gt11. *Proc Natl Acad Sci USA* 1987; 84: 5350-4.
- Paxton RJ, Mooser G, Pande H, Lee TD, Shively JE. Sequence analysis of carcinoembryonic antigen: identification of glycosylation sites and homology with the immunoglobulin supergene family. *Proc Natl Acad Sci USA* 1987; 84: 920-4.
- Zimmermann W, Weber B, Ortlieb B, Rudert F, Schempp W, Fiebig HH, Shively JE, Von Kleist S, Thompson JA. Chromosomal localization of the carcinoembryonic antigen gene family and differential expression in various tumors. *Cancer Res* 1988; 48: 2550-4.
- Thompson JA, Mauch EM, Chen F-C, Hinoda Y, Schrewe H, Berling B, Barnert S, von Kleist S, Shively JE, Zimmermann W. Analysis of the size of the carcinoembryonic antigen (CEA) gene family: isolation and sequencing of N-terminal domain exons. *Biochem Biophys Res Comm* 1989; 158: 996-1004.
- Thompson J, Koumari R, Wagner K, Barnett S, Schleussner C, Schrewe H, Zimmermann W, Muller G, Schempp W, Zaninetta D, Ammaturo D, Hardman N. The human pregnancy-specific glycoprotein genes are tightly linked on the long arm of chromosome 19 and are coordinately expressed. *Biochem Biophys Res Comm* 1990; 167: 848-59.
- Oikawa S, Imajo S, Noguchi T, Kosaki G, and Nakazato, H. The carcinoembryonic antigen (CEA) contains multiple immunoglobulin-like domains. *Biochem Biophys Res Comm* 1987; 142: 634-42.
- Williams AF, Barclay AN. The immunoglobulin superfamily: domains for cell surface recognition. *Ann Rev Immunol* 1988; 6: 381-405.
- Hefta SA, Hefta LJF, Lee TD, Paxton RJ, Shively JE. Carcinoembryonic antigen is anchored to membranes by covalent attachment to a glycosylphosphatidylinositol moiety: identification of the ethanolamine linkage site. *Proc Natl Acad Sci USA* 1988; 85: 4648-52.
- Takami N, Misumi Y, Kuroki M, Matsuoka Y, Ikehara Y. Evidence for carboxyl-terminal processing and glycolipid-anchoring of human carcinoembryonic antigen. *J Biol Chem* 1988; 263: 12716-20.
- Kolbinger F, Schwarz K, Brombacher F, Von Kleist S, Grunert F. Expression of an NCA cDNA in NIH/3T3 cells yields a 110K glycoprotein, which is anchored into the membrane via glycosyl-phosphatidylinositol. *Biochem Biophys Res Comm* 1989; 161: 1126-34.
- Berling B, Kolbinger F, Grunert F, Thompson JA, Brombacher F, Buchegger F, von Kleist S, Zimmermann W. Cloning of a carcinoembryonic antigen gene family member expressed in leukocytes of chronic myeloid leukemia patients and bone marrow. *Cancer Res* 1990; 50: 6534-9.
- Hinoda Y, Neumaier M, Hefta SA, Drzeniek Z, Wagener C, Shively L, Hefta LJF, Shively JE, Paxton RJ. Molecular cloning of a cDNA coding biliary glycoprotein I: primary structure of a glycoprotein immunologically crossreactive with carcinoembryonic antigen. *Proc Natl Acad Sci USA* 1988; 85: 6959-63.
- Barnett T, Kretschmer A, Austen DA, Goebel SJ, Hart JT, Elting JJ, Kamarck ME. Carcinoembryonic antigens: alternative splicing accounts for the multiple mRNAs that code for novel members of the carcinoembryonic antigen family. *J Cell Biol* 1989; 108: 267-76.
- Benchimol S, Fuks A, Jothy S, Beauchemin N, Shiota K, Stanners CP. Carcinoembryonic antigen, a human tumor marker, functions as an intercellular adhesion molecule. *Cell* 1989; 57: 27-34.
- Oikawa S, Inuzuka C, Kuroki M, Matsuoka Y, Kosaki

- G, Nakazato H. Cell adhesion activity of non-specific cross-reacting antigen (NCA) and carcinoembryonic antigen (CEA) expressed on CHO cell surface: homophilic and heterophilic adhesion. *Biochem Biophys Res Comm* 1989; 164: 39-45.
22. Oikawa S, Inuzuka C, Kuroki M, Arakawa F, Matsuo Y, Kosaki G, Nakazato H. A specific heterotypic cell adhesion activity between members of carcinoembryonic antigen family, W272 and NCA, is mediated by N-domains. *J Biol Chem* 1991; 266: 7995-8001.
23. Rojas M, Fuks A, Stanners CP. Biliary glycoprotein, a member of the immunoglobulin supergene family, functions *in vitro* as a  $\text{Ca}^{2+}$ -dependent intercellular adhesion molecule. *Cell Growth and Diff* 1990; 1: 527-33.
24. Zhou H, Fuks A, Stanners CP. Specificity of intercellular adhesion mediated by various members of the immunoglobulin supergene family. *Cell Growth and Diff* 1990; 1: 209-15.
25. Leusch HG, Hefta SA, Drzeniek Z, Hummel K, Markos-Pusztai Z, Wagener C. *Escherichia coli* of human origin binds to carcinoembryonic antigen (CEA) and non-specific crossreacting antigen (NCA). *FEBS Letters* 1990; 261: 405-9.
26. Toth CA, Rapoza A, Kowal A, Steele GD, Thomas P. Receptor-mediated endocytosis by human Kupffer cells. *Biochem Soc Trans* 1988; 16, 1027-8.
27. Takeichi M. Cadherin cell adhesion receptors as a morphogenetic regulator. *Science* 1991; 251, 1541-55.
28. Albelda SM, Muller WA, Buck CA, Newman PJ. Molecular and cellular properties of PECAM-1 (endoCAM, CD 31): a novel vascular all-cell adhesion molecule. *J Cell Biol* 1991; 114, 1059-68.
29. Lin SH, Fain JN. Purification of  $(\text{Ca}^{2+}$ - $\text{Mg}^{2+}$ )-ATPase from rat liver plasma membranes. *J Biol Chem* 1984; 259: 3016-20.
30. Obrink B. C-CAM (Cell-CAM 105) - A member of the growing immunoglobulin superfamily of cell adhesion proteins. *Bioessays* 1991; 13: 227-34.
31. Lin SH, Guidotti G. Cloning and expression of a cDNA coding for a rat liver plasma membrane ecto-ATPase: the primary structure of the ecto-ATPase is similar to that of the human biliary glycoprotein I. *J Biol Chem* 1989; 264: 14408-14.
32. Pollard JW, Stanners CP. Characterization of cell lines showing growth control isolated from both the wild type and leucyl-tRNA synthetase mutant of Chinese hamster ovary cells. *J Cell Physiol* 1979; 98: 571-85.
33. Turbide C, Rojas M, Stanners CP, Beauchemin N. A mouse carcinoembryonic antigen gene family member is a calcium-dependent cell adhesion molecule. *J Biol Chem* 1991; 266: 309-15.
34. McCuaig K, Turbide C, Beauchemin N. mmCGM1a: a mouse carcinoembryonic antigen gene family member, generated by alternative splicing, functions as an adhesion molecule. *Cell Growth and Diff* 1992; 3.
35. Hasegawa T, Isobe K, Tsuchiya Y, Oikawa S, Nakazato H, Ikezawa H, Nakashima I, Shimokata K. Establishment and characterization of human carcinoembryonic antigen transgenic mice. *Br J Cancer* 1991; 64, 710-4.